

On a possible dynamical scenario leading to a generalised Gamma distribution

Silvio M. Duarte Queirós*

Centro Brasileiro de Pesquisas Físicas, 150, 22290-180 Rio de Janeiro-RJ, Brasil

(February 2, 2008)

In this report I present a possible scenario which can lead to the emergence of a generalised Gamma distribution first presented by C Tsallis *et al.* as the distribution of traded volumes of stocks in financial markets. This propose is related with superstatistics and the notion of moving average commonly used in econometrics.

I. THE Γ -DISTRIBUTION

The Γ -distribution is a general distribution that is verified in processes where the waiting times between variables that follow a Poisson distribution are significant. It involves two free parameters, usually labeled by α and θ and defined as [1],

$$p_{\alpha,\theta}(x) = \frac{x^{\alpha-1} \exp\left(-\frac{x}{\theta}\right)}{\Gamma[\alpha] \theta^\alpha}. \quad (1)$$

A special case of Γ -distribution is to consider $\alpha = r/2$ and $\theta = 2$. In this case the distribution is called χ^2 -distribution and represents the probability of get a value ζ of a variable that is obtained by the summation of r independent squared variables ξ_i associated with the Gaussian distribution with null mean and unitary variance [1],

$$\zeta = \sum_{i=1}^r \xi_i^2. \quad (2)$$

The same form presented in Eq. (1) can be obtained as the stationary solution of the following differential stochastic equation

*Electronic address: sdqueiro@cbpf.br

$$dx_t = -\gamma (x_t - \theta) dt + k\sqrt{x}dW_t. \quad (3)$$

Considering the Itô convention for stochastic differentials I am able to write the Fokker-Planck equation [2],

$$\frac{\partial p(x, t)}{\partial t} = \frac{\partial}{\partial x} [\gamma (x_t - \theta) p(x, t)] + \frac{1}{2} \frac{\partial^2}{\partial x^2} [k^2 x p(x, t)], \quad (4)$$

whose stationary solution is

$$p_{\alpha, \theta}(x) = \frac{\alpha^\alpha}{\Gamma[\alpha]} \frac{x^{\alpha-1}}{\theta^\alpha} \exp\left(-\frac{\alpha}{\theta}x\right) \quad (5)$$

with $\alpha = 2\gamma\theta k^{-2}$. Performing a simple variable change $x \rightarrow \frac{x}{\alpha}$, it is possible to transform Eq. (5) into Eq. (1) and the Itô-Langevin equation (3)

$$dx = -\gamma (x - \alpha\theta) dt + \sqrt{2\gamma\theta}\sqrt{x} dW_t. \quad (6)$$

For a question of simplicity let me represent θ as β^{-1} . So Eq. (1) will be written as,

$$p_{\alpha, \beta}(x) = \frac{\beta^\alpha}{\Gamma[\alpha]} x^{\alpha-1} \exp(-\beta x). \quad (7)$$

In figure 1 are depicted some examples of $\Gamma(x)$ distributions.




FIG. 1. Representation of $\Gamma(x)$ vs x for some values of α and θ . (I) - $\alpha = 2$ and $\theta = 2$; (II) - $\alpha = 4$ and $\theta = 2$; (III) - $\alpha = 2$ and $\theta = 4$. The inset presents the same curves, but in a log-log scale.

II. INTRODUCING THE GENERALISED Γ -DISTRIBUTION

Let one now suppose that parameter θ in Eq. (3) is in fact a stochastic variable on time scale larger than the characteristic time scale γ^{-1} . This means that $p_{\alpha,\beta}(x)$ is, for this case, a conditional probability density function $p_{\alpha}(x|\beta)$. If the random process for β is associated with a SPDF, $\Pi(\beta)$, then the SPDF for x variable, $P(x)$, is simply given by

$$P(x) = \int p_{\alpha}(x|\beta) \Pi(\beta) d\beta. \quad (8)$$

Among the various distributions for non-negative variables let one consider that β is associated, itself, with a Γ -distribution,

$$P(x) = \int p_{\alpha}(x|\beta) \Pi(\beta) d\beta. \quad (9)$$

which can be associated to a microscopic equation similar to Eq. (6).

Calculating the integral presented in equation (8) one gets,

$$P(x) = \frac{\Gamma[\alpha + \lambda] \omega^{\alpha}}{\Gamma[\alpha] \Gamma[\lambda]} x^{\alpha-1} (1 + \omega x)^{-(\alpha+\lambda)}. \quad (10)$$

Defining $\bar{\theta} = \frac{1}{\omega(\alpha+\lambda)}$ and $q = 1 + \frac{1}{\alpha+\lambda}$, Eq. (10) can be rewritten as,

$$P(x) = \frac{\Gamma\left[\frac{1}{q-1}\right] \left(\frac{q-1}{\bar{\theta}}\right)^{\alpha}}{\Gamma[\alpha] \Gamma\left[\frac{1}{q-1} - \alpha\right]} x^{\alpha-1} \left\{1 - (1-q) \frac{x}{\bar{\theta}}\right\}^{\frac{1}{1-q}}, \quad (11)$$

$$P(x) \equiv \frac{\Gamma\left[\frac{1}{q-1}\right] \left(\frac{q-1}{\bar{\theta}}\right)^{\alpha}}{\Gamma[\alpha] \Gamma\left[\frac{1}{q-1} - \alpha\right]} x^{\alpha-1} \exp_q\left[-\frac{x}{\bar{\theta}}\right], \quad (12)$$

which I will call the ${}_q\Gamma$ -distribution. This kind of distribution was already verified, at least, for the distribution of traded volumes of stocks in financial markets [3]. For the limit $q \rightarrow 1$, the usual Γ -distribution is recovered, which corresponds to $\Pi(\beta) = \delta\left(\beta - \frac{1}{\bar{\theta}}\right)$. Some examples of ${}_q\Gamma$ -distribution are presented in Fig. 2,




FIG. 2. Representation of $\Gamma(x)$ vs x for some values of α and θ . (I) - $q = 1.2$, $\alpha = 2$ and $\theta = 2$; (II) - $q = 1.1$, $\alpha = 2$ and $\theta = 2$; (III) - $q = 1.1$, $\alpha = 2$ and $\theta = 4$; (IV) - $q = 1.1$, $\alpha = 4$ and $\theta = 2$. The inset presents the same curves, but in a log – log scale.

. The same form is presented in Figs. 7 and 8 of Ref. [3].

This problem of fluctuations in some intensive parameter of the dynamical equation(s) that describe(s) the evolution of a system [4] was recently studied by C. Beck in the context of Langevin equation with fluctuating temperature [5] and extended together with Eddie G.D. Cohen [6] who defined it as *superstatistics* (a statistic of statistics). This superstatistics presents a close relation to the non-extensive statistical mechanics framework based on the entropic form [7,8],

$$S_q = \frac{1 - \int [p(x)]^q dx}{q - 1}. \quad (13)$$

For the problem of the distribution of traded volume of stocks in financial markets the presence of fluctuations in θ , or the mean value of the scaled variable αx , it is similar to the problem of the moving average in the analysis of the volatility useful in the reproduction of some empirical facts like the autocorrelation function and the so-called leverage effect, see e.g. Ref. [9].

- [1] M.V. Jambunathan, Ann. Math. Stat. 25 (1954) 401;
- [2] H. Risken, The Fokker-Planck Equation - Methods of Solution and its Applications, 2nd edition, (Springer-Verlag, Berlin) 1989;
- [3] R. Osório, L. Borland and C. Tsallis, *Distributions of High-Frequency Stock-Market Observables* in; M. Gell-Mann and C. Tsallis, *Nonextensive Entropy - Interdisciplinary Applications* (Oxford University Press, New York, 2004);
- [4] G. Wilk and Z. Włodarczyk, Phys. Rev. Lett. 84 (2000) 2770;
- [5] C. Beck, Phys. Rev. Lett. 87 (2001) 180601;
- [6] C. Beck and E.G.D. Cohen, Physica A 322 (2003)267;
- [7] C. Tsallis, J. Stat. Phys. **52** , 479 (1988). A regularly an updated bibliograhya on the subject is available at <http://tsallis.cat.cbpf.br/biblio.htm>;
- [8] M. Gell-Mann and C. Tsallis, *Nonextensive Entropy - Interdisciplinary Applications* (Oxford University Press, New York, 2004);
- [9] J. Perelló, J. Masoliver and J.P. Bouchaud, cond-mat/0302095 (preprint, 2003).